

*Funtastic Fluoride...*

# PediaGel<sup>®</sup>

**TOPICAL FLUORIDE GEL**

**CAUTION:**

U. S. Federal Law prohibits dispensing without prescription.

**DESCRIPTION:**

PediaGel Acidulated Phosphate Fluoride Topical Gel is for professional topical application as a fluoride treatment. Contains 1.23% acidulated phosphate fluoride [1.23% fluoride ion per each gram or milliliter]. For MSDS visit [preventech.com](http://preventech.com).

**INDICATIONS AND USAGE:**

For topical application to aid in the protection against dental caries.

**CONTRAINDICATIONS:**

Hypersensitivity to fluoride.

**WARNINGS:**

Do not swallow. If treatment dose is swallowed [less than 100 mg F], administer milk, limewater, or calcium-type antacid. In case of larger dose use ipecac syrup emetic and immediately seek medical help. May contain: FD&C Red 40, D&C Red 33, and/or FD&C Blue 1.

**PRECAUTIONS:**

For dental professional use only. Safety and effectiveness below age 3 have not been established. There have been no long-term animal studies with this product to evaluate cariogenic, mutagenic, or impairment of fertility potential. Laboratory studies have indicated that repeated use of APF may dull porcelain, composite restorations, and sealants.

**ADVERSE REACTIONS:**

Developing teeth of children under age 6 may become permanently discolored if excessive amounts are repeatedly swallowed. The following adverse reactions are possible in individuals hypersensitive to fluoride: eczema, atropic dermatitis, urticaria, gastric distress, headache, and weakness.

**DOSAGE and ADMINISTRATION:**

- 1) Remove cap and induction seal. Replace cap.
- 2) Dispense gel into foam application tray.
- 3) Air dry teeth thoroughly and insert tray in mouth with head slightly tilted forward.
- 4) Instruct patient to continue light biting action for 1 minute (or up to 4 minutes). A slight chewing motion enhances interproximal coverage.
- 5) Use suction throughout treatment.
- 6) Have patient expectorate after treatment.
- 7) Instruct patient not to drink or rinse for 15 minutes or eat for 30 minutes.

**RECOMMENDED FREQUENCY:**

Not to exceed 4 treatments per year.

**HOW SUPPLIED:**

12 Fl. oz (345mL)

**STORE AT ROOM TEMPERATURE**

**PROTECT FROM FREEZING**

**KEEP OUT OF REACH OF CHILDREN**

**R<sub>x</sub>only**

PREVENTECH<sup>®</sup>



4330-C Matthews Indian Trail Road • Indian Trail, NC 28079

800.474.8681 • 704.849.2416 • fax: 704.849.2417 • [preventech.com](http://preventech.com)

